

Homework 4: Life History Evolution (Updated)

1. Invasion fitness

In class we used an adaptive dynamics approach to study the evolution of carrying capacity in the logistic model K . Let's do the same for demographic traits in the Beverton-Holt model of growth. From homework 2 we have that the population dynamics in the Beverton-Holt model are given by:

$$N(t+1) = \lambda N(t) \frac{1}{1 + \alpha N(t)}$$

- Start by assuming the $\alpha(z) = z$. Use an adaptive dynamics approach to determine whether evolution should favour an increase or decrease in α .
- Repeat part A but now for other demographic trait λ (assuming α is constant).
- Propose functions $\alpha(z)$ and $\lambda(z)$ that may give you a finite evolutionary singular strategy. You need not show that it does.

2. Mutation Accumulation

In class we considered the evolution of an antagonistic-pleiotropic allele that have fitness benefits and young ages and deleterious effects at older ages. A non-mutually exclusive explanation for the evolution of senescence is that deleterious mutations are under less selection at older ages and hence obtain higher frequencies.

- Propose a model of mutation-selection balance in a non-age structured population. Consider a life cycle of Census->Mutation->Selection->Reproduction. Assume that the mutation is deleterious

$$W_{AA} = (1 - 2s), W_{Aa} = (1 - 2s), \text{ and } W_{aa} = 1$$

and that mutations from A->a and a->A occur at the same rate μ .

What is the equilibrium allele frequency of the mutation and how does it depend on s and μ ?

- Suppose that in the absence of mutation the death rate is d_0 with the probability of survival being $M = 1 - d_0$. The mutation increases death rate.

$$d_{AA} = d_0(1 + 2s), d_{Aa} = d_0(1 + 2s) \text{ and } d_{aa}(a) = d_0$$

Consider a model with three age classes and an age-independent fecundity of F . Calculate the expression for the quasi-equilibrium and young-only approximations for the fitness of a mutant. Compare these fitnesses to the analogous fitnesses in the wild-type population.

- Analyzing a Leslie matrix numerically (assume $d_0 = 0.3$, $F = 1$, $s = 0.1$) compare the stable age distribution of the mutant and the wild-type.

3. Seed Dormancy with seedling survival

In perennials newly germinated seeds have a different probability of survival than adult plants (e.g., raspberries). To incorporate this into our perennial model of seed dormancy evolution let's introduce an environmental-dependent juvenile survival s_{Ji} .

- Propose a recursion equation for this system.
- Calculate the long-term growth rate. Characterize numerically under what conditions does the population grow? Assume $\bar{s}_J < \bar{s}_A$.
- Use a combination of analytical and numerical methods to explore how juvenile survival impacts the evolution of seed dormancy. You will have to make assumptions about correlations between environmental-dependent components– justify your choices.

